



The phenomenon of bioluminescence in insects

Alaa H. Oleiwi¹, Sukania R. Neamah² and Ali M. Atshan²

¹Department of Biology, College of Sciences, University of Thi-Qar

²General Directorate of Education in Thi-Qar province

*Corresponding author: ala.oleiwi@sci.utq.edu.iq, sukainarn2@gmail.com, msahir21@Yahoo.com

Received: Feb 12, 2021 / Revised Mar 12, 2021/ Accepted: Mar 26, 2020

Abstract

The light emitted by many insect species of insects and other organisms are known as bioluminescence. Dating back to the earliest written records of the ancient Greeks, bioluminescence is one of the ancient fields of scientific study. There are many studies on insects mentioned in this article. In the activities of insects, bioluminescence plays a significant function.

Keywords Luciferases, bioluminescence, Insects, importance of bioluminescence.

Introduction

Bioluminescence is a biological process that usually occurs in the process of transferring energy in the form of light through a synthetic response. Life forms such as microscopic organisms, growths, mushrooms, phytoplankton, marine protists, limb animals, mollusks, cents, millipedes, shellfish, and chords show light. Besides, some novel creepy reptiles can also show light. Although bioluminescence is created by different creatures, it has evolved abnormally in bed bugs, and bed bugs play a key role in romance, predation, and protection (Sankarganesh and Rajeshwaran, 2019).

Bioluminescence Origin

At the point where species produce and radiate light as a result of a compound reaction, the cycle is called bioluminescence-organism means "life" in Greek, and "lumen" means "light" in Latin. The composite energy is converted to light energy throughout the cycle (Longkumer and Kumar, 2018).

The phenomenon of bioluminescence

This phenomenon occurs when there is a pigment called Lucifer and an enzyme called luciferase with a source of oxygen. As a result of the reaction, a compound called (oxyluciferin) is produced and emits light (Fig. 1) Luciferase is a key compound that catalyzes the luminescence reaction in bioluminescence. Scheme (1) (Longkumer and Kumar, 2018).





Fig. 1 Structure of Luciferin

They catalyze the oxygenation of the mixture traditionally called luciferin and produce energy-rich peroxide intermediates. Their unlimited degradation will produce singlet electron-excited substances, which will decay with high efficiency and produce obvious Photon photon (Viviani, 2002; Vojtek, *et al.*, 2014)

Bioluminescence in some orders of Insects

Coleoptera (beetles)

The beetle has the largest number of luminous species and the largest variety. It also includes fireflies (Lampyridae), railroad worms (Phenodidae), and click bugs (Elateridae). Luminescence was also found in *Xantholinus* larvae of (Staphylinidae) (Costa, et al., 1986).

Fireflies will produce green-yellow flames from the length, span, and recurrence of the belly lamp for sexual purposes. Bug insects generally emit a continuous green light, and the abdominal light organ in the abdomen emits a continuous green-orange light. Railroad worms transmit the largest color among glow beetles, including the most explosive larvae, while females are cylindrical. Horizontal lights along the body produce green-orange light.

Fireflies produce yellow and green flames, and the length, span, and lights emitted from the abdomen of these flames are used for sexual purposes. When soaring, the organs on the abdominal belly of a creeping reptile emit a continuous green-orange light, while bugs emit continuous green light. The most important colors of glow beetles are transmitted by railroad worms, including the most explosive larvae and females.

A horizontal column of lights along the body produces greenish-orange light (Seliger, et al., 1964; Moiseff, et al., 1999; Bechara, 1988).

Diptera

Mycetophilidae is one of the most widespread families that have luminescence which includes the most common species of *Arachnocampa* found in Australian and New Zealand caves, and whose young are found at the top of the caves with the network (Meyer, 1990).

The light organs in the five anterior segments and the five anterior segments and the tail are located and are composed of black bodies, which are filled with dark particles secreted by huge cellular mitochondria (Bassot, 1978)

Hemiptera

There are strange and attractive insects in the lantern fly (Fulgoridae). Many Saxifrage species are brightly colored, and some of the produced epidermal wax is mainly composed of keto Esters (Mason et al., 1989).

Advantages of bioluminescence in Insects

Defense

The constant glow of the head in railroad worm larvae while walking suggests that it may have an illumination role, whereas the condition of turning on the sidelight organs indicates a defensive function. Potential predators may be frightened away by the unexpected flash. (Longkumer and Kumar, 2018).

Predation

The most special example of illumination is light acting. In New Zealand, the urge to hunt for prey is powerful. *Arachnocampa Luminosa*, or fireflies, fly. Female flies lay their eggs in the cave with the low ceiling. The larva stays down after hatching. Glue the thread in place and watch it shine. This light can glow in the cave at night, attracting other insects. Insects are trapped in sticky strings and are preyed upon by larvae. Cave People who live in the flies are generally called "luminous caves" and are tourist attractions in New Zealand (Longkumer and Kumar, 2018).

Mating signal

It's believed the fireflies use light as a mating signal. Bioluminescence attracts members of the same group to congregate, increasing the likelihood of mating in certain animals. Female *Chlorophytum* with no wings, sedentary in some organisms (Longkumer and Kumar, 2018).

Previous study about the phenomenon of bioluminescence

The majority of the research on insect luciferases and biochemistry has been done with fireflies. The bulk of research on insect luciferases and biochemistry has been conducted on fireflies. The study revealed that substrate deflation is a prevalent trend in this class of legacies. The three-dimensional structure of firefly luciferase was recently identified. (Viviani, 2002).

The first study of De Cock and Matthysen's (2003) Lampyrid larvae used luminescence to alert nocturnal, visual predators that they are dangerous. Attack latencies for luminescent prey rose after exposure to glow-worm eggs, which were considered unpleasant by the toads, but not for non-glowing prey.

Bioluminescence is one of the oldest fields of scientific inquiry, dating back to the first written documents of the ancient Greeks. This article discusses the various animal luminescence experiments carried out until the end of the 19th century (Lee, 2008). Another finding by Costa and Vanin (2010) that was carried out in Central Brazil's bioluminescent termite nests and for this purpose, general aspects of elaterid fireflies-related bioluminescence were also given. Adaptations for all beetle larvae to live in the nest setting and practical association groups.

(Vojtek et al., 2014) Whereas the other study was using all bacterial species and *Bombyx Mori* or *Galleria mellonella* haemolymph, we observed a large dose-dependent decrease in bioluminescence, which was confirmed by a drop in bacterial viability as measured by plating. The humoral origin of the antibacterial activity found in total hemolymph was confirmed in haemolymph plasma lacking haemocytes.

Despite the widespread biotechnological applications of firefly luciferase and the evolution of firefly flash signals, firefly bioluminescence has been observed in the Big Dipper firefly *Photinus pyralis*. The main research center luciferase

was cloned from the fly from which the main research centre luciferases were cloning (Fallon *et al.*, 2017).

It is understood that light in fireflies serves as a mating signal. In certain animals, bioluminescence attracts members of the same group to congregate, thus increasing the chance of mating. wingless female Chlorophytum, sedentary in some animals. (Longkumer and Kumar, 2018).

(Owens and Lewis 2018) which Long-wavelength ambient light had little effect on signal morphology, owing. These findings suggest that long-wavelength lighting has less of an effect on firefly courtship than broad-spectrum white lighting, it's also possible that using it instead of wide-spectrum white lighting could help firefly conservation efforts.

Another study by (Adams and Miller 2019) discovered Fireflies, a species of beetle that uses D-luciferin as a substrate, the most famous examples of luminescence. The luminescence that occurs in the beetles arises from the fatty acyl-CoA synthetase enzymes that are generally found in insects. ACS replication and subsequent diversification were the drivers of the evolution of the Bioluminescence, according to the recent sequence of firefly genomes and transcription. Bioluminescence is produced by different species and is highly evolved in insects that play a major role in courtship, predation, and defense.

Conclusions

Luciferase is a major compound that catalyzes the bioluminescence scintillation reaction. It stimulates the oxygenation of the mixture conventionally called luciferin and produces an energy-rich peroxide medium.

Bioluminescence was found in some orders of insects and have several functions, either as a defence, mating and predation. Bioluminescence have been studied by a number of researchers from many aspects.

Conflict of Interest

The author hereby declares no conflict of interest.

Consent for publication

The author declares that the work has consent for publication.

Funding support

The author declares that they have no funding support for this study.

References

- Adams Jr, ST, & Miller, SC (2020). Enzymatic promiscuity and the evolution of bioluminescence. *The FEBS journal*, 287 (7), 1369-1380.
- Bassot, J. M., & JM, B. (1978). Les corps noirs, cellules géantes du Diptère mycétophilide lumineux *Platyura fultoni* et leur sécrétion mitochondriale. *C. R. Acad. Sci. Paris* 287: 623–630.
- Bechara, E. J. (1988). Luminescent elaterid beetles: biochemical, biological and ecological aspects. *Advances in oxygenated processes*, 1, 123-178.
- Costa, C., Vanin, S. A., & Neto, P. C. (1986). Larvae of neotropical Coleoptera. XIV: First record of bioluminescence in the Family Staphylinidae (Xantholini). *Revista brasileira de entomologia*, 30(1), 101-104.
- Costa, C., & Vanin, S. A. (2010). Coleoptera larval fauna associated with termite nests (Isoptera) with emphasis on the “bioluminescent termite nests” from Central Brazil. *Psyche*, 2010.
- De Cock, R., & Matthysen, E. (2003). Glow-worm larvae bioluminescence (Coleoptera: Lampyridae) operates as an aposomatic signal upon toads (*Bufo bufo*). *Behavioral Ecology*, 14(1), 103-108.
- Fallon, T. R., Lower, S. E., Chang, C. H., Bessho-Uehara, M., Martin, G. J., Bewick, A. J., ... & Weng, J. K. (2018). Firefly genomes illuminate parallel origins of bioluminescence in beetles. *Elife*, 7, e36495. <https://doi.org/10.1101/237586>.
- Lee, J. (2008). Bioluminescence: the First 3000 Years (Review). *Journal of Siberian Federal University. Biology* 3 (194-205).
- Longkumer, I. Y., & Kumar, R. (2018). Bioluminescence in Insect. *Int. J. Curr. Microbiol. App. Sci*, 7(3), 187-193.
- Mason, R. T., Fales, H. M., Jones, T. H., O'Brien, L. B., Taylor, T. W., Hogue, C. L., & Blum, M. S. (1989). Characterization of fulgorid waxes (homoptera: fulgoridae: insecta). *Insect biochemistry*, 19(8), 737-740.
- Meyer-Rochow V. B. (1990). The New Zealand Glow-Worm. Waitomo Caves Museum Society, Waitomo Caves, New Zealand.
- Moiseff, A., Metcalfe, J., Copeland, J., & Palmieri, F. (1999). Synchrony in fireflies: Diversity, mechanisms, and model. *Bioluminescence and chemiluminescence: Perspectives for the 21st century*, 573-576.
- Sankarganesh, E. and Rajeshwaran, B. (2019). Bioluminescence insects and its significance in entomology. *AGRICULTURE & FOOD*; Volume 1 – Issue 12 – December 2019.
- Seliger, H. H., Buck, J. B., Fastie, W. G., & McElroy, W. D. (1964). The spectral distribution of firefly light. *The Journal of general physiology*, 48(1), 95-104.
- Viviani, V. R. (2002). The origin, diversity, and structure function relationships of insect luciferases. *Cellular and Molecular Life Sciences CMLS*, 59(11), 1833-1850.
- Vojtek, L., Dobeš, P., Büyükgüzel, E., Atosuo, J., & Hyršl, P. (2014). Bioluminescent assay for evaluating antimicrobial activity in insect haemolymph. *Eur. J. Entomol.* 111(3): 335–340.
- Owens, A. C., & Lewis, S. M. (2018). The impact of artificial light at night on nocturnal insects: A review and synthesis. *Ecology and evolution*, 8(22), 11337-11358.

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).



How to cite this article:

Oleiwi, A. H., Neamah, S. R. and Atshan, A. M. (2021). The phenomenon of Bioluminescence in Insects. *Science Archives*, Vol. 2 (1), 62-64. <http://dx.doi.org/10.47587/SA.2021.2110>