



A systematic review of tapeworms (Plathelminthes, Cestoda) of *Columba livia*

Nuha Jabbar Alrikaby¹ and Hasan Azaati Ibadi Abul-Doanej²

¹Department of Biology, College of Science, University of Thi Qar, Iraq

²Department of specialist supervision/Directorate General of Mysan Education/Ministry of Education, Iraq

Received: Apr 7, 2023/ Revised: May 4, 2023/ Accepted: May 6, 2023

(✉) Corresponding Author: Nuha_bio@sci.utq.edu.iq

Abstract

After conducting research and analyzing literature data, it was discovered that Iraq is home to tapeworms belonging to four distinct families, namely Davaineidae, Hymenolepididae, Dilepididae, and Anoplocephalidae. Within these families, several species of tapeworm have been identified in Iraq, including *Cotugnia intermedia*, *Raillietina tetragona*, *Raillietina cesticillus*, *Raillietina echinobothrida*, *Aporina delafondi*, and *Davainea proglottina*. *Cotugnia intermedia* is a tapeworm commonly found in the intestines of small rodents and is transmitted to cats and dogs through the ingestion of infected rodents. *Raillietina tetragona* and *Raillietina cesticillus* are two species of tapeworm that commonly infect chickens, while *Raillietina echinobothrida* is found in pigeons and doves. *Aporina delafondi* is a tapeworm that has been found in the intestines of cattle, sheep, and goats in Iraq. The adult worms attach to the host's intestinal wall and can cause intestinal inflammation and blockages. Finally, *Davainea proglottina* is a tapeworm that infects the intestines of domestic and wild carnivores, including dogs, cats, and foxes. Infection occurs through ingestion of infected intermediate hosts such as rodents, birds, and reptiles. Identifying these tapeworm species in Iraq is important for developing effective control and prevention strategies, as many of these parasites can cause significant health problems in animals and humans. Proper sanitation and hygiene practices, as well as deworming programs for domestic animals, are important measures that can help to reduce the spread of these tapeworms and their associated diseases.

Keywords: Plathelminthes, Cestoda, *Columba livia*, Species identification, Control strategies, Health problems.

Introduction

The tapeworms are a group of endoparasitic helminths that almost exclusively occupy their definitive host's alimentary canal (Smyth and McManus, 2007).

Pigeons (order Columbiformes) are widespread and can be found in every town and city around the world (Marques et al., 2007). The effects of parasitism on birds are often serious (Dranzoa, 1999).

Many helminthological studies devoted to studying the parasite fauna including cestodes of domestic pigeons have been carried out in many countries in India (Magzoub et al., 1980; in India Malhotra and Kapoor, 1983; Adang et al., 2008) in Thessaloniki; Northern Greece (Diakou et al., 2013)

northern Thailand (Butboonchoo, et al., 2016) as well as in Iraq (Al-Bayati, 2011) in Iraq (Hassan, 2013; Al-Marsomy and Al-Hamadaani, 2016; Abed Alrikaby et al., 2020 and Al-Hussaini and Afrah, 2021).

It should be noted that all the above studies were including identification, biochemistries, and histopathological studies and there is no review information on domestic pigeons.

The present review aims to epitomize the information on cestode parasites of *Columba livia* which have been recorded worldwide until 2021.

The arrangement of the cestoda follow the classification and nomenclature by Khalil et al. (1994) with updates by Caira et al. (2014).

Phylum: Platyhelminthes ([Gegenbaur, 1859](#))
Class: Cestoda ([Rudolphi, 1808](#))
Order: Cyclophyllidea ([Braun, 1900](#))
Family: Davaineidae ([Braun, 1900](#))
Genus: Cotugnia ([Diamare, 1893](#))
Cotugnia digonopora ([Pasquale, 1890](#))
Intermediate hosts: insects
Habitat: small intestine
Distribution: in India ([Waghmare et al., 2009](#)).

Cotugnia intermedia

Intermediate hosts: Habitat
Distribution in Iraq ([Al-Bayati, 2011](#)).

Cotugnia polycantha

Intermediate hosts: insects
Habitat: small intestine
Distribution: Egypt ([Ahmed, 2012](#))

Cotugnia cuneata in Bangladesh ([Musa et al., 2011](#))

Intermediate hosts
Habitat small intestine
Distribution

Cotugnia satpuliensis

Intermediate hosts
Habitat small intestine
Distribution in India ([Malhotra and Kapoor, 1983](#))

Cotugnia srivastavai

In India ([Malviya and Dutt, 1970](#)).
Intermediate hosts
Habitat small intestine
Distribution

Cotugnia majdoubii in India ([Magzoub et al., 1980](#)).

Phylum: Platyhelminthes ([Gegenbaur, 1859](#))

Class: Cestoda ([Rudolphi, 1808](#))
Order: Cyclophyllidea ([Braun, 1900](#))
Family: Davaineidae ([Braun, 1900](#))
Genus: *Raillietina* ([Fuhrmann, 1920](#))
Raillietinia canabia in India ([Magzoub et al., 1980](#))
Raillietinia zahratis in India ([Magzoub et al., 1980](#))
Raillietina echinobothrida ([Megnin, 1881](#))
Intermediate hosts: insects
Habitat: small intestine Duodenum/ileum
Distribution: Cosmopolitan. Thessaloniki; Northern Greece ([Diakou et al., 2013](#))

Raillietina tetragona

Intermediate hosts: insects ant
Habitat: small intestine Duodenum/ileum
Distribution: Cosmopolitan. Thessaloniki; Northern Greece ([Diakou et al., 2013](#)) ([Adang et al., 2008](#)) (in Thailand [Butboonchoo, et al., 2016](#)).

Raillietina cesticillus

Intermediate hosts: beetle
Habitat: small intestine Ileum
Distribution: Cosmopolitan. Thessaloniki; Northern Greece ([Diakou et al., 2013](#)) (Nigeria [Adang et al., 2008](#)) ([Hassan, 2013](#) in Iraq) (in Iran [Radfar, et al 2012](#)) (in Thailand [Butboonchoo, et al., 2016](#)).

Raillietina micracantha

Intermediate hosts: insects
Habitat: small intestine
Distribution: Cosmopolitan Spain ([Torres et al., 2010](#))

Raillietina vinagoi

Intermediate hosts: insects
Habitat: small intestine
Distribution: (in, Egypt [Ghobashy and Taeleb, 2015](#)).

Raillietina magninumida

Intermediate host: beetles
Habitat: small intestine
Distribution ([Radfar, et al., 2012](#)).

Raillietina echinobothrida

Intermediate hosts: ants
Habitat: small intestine
Distribution: (in Iraq [Al-Marsomy and Al-Hamadaani, 2016](#)). in Bangladesh [Musa et al., 2011](#) in Bangladesh) (Nigeria [Adang et al., 2008](#)) (in Iraq [Al-Marsomy and Al-Hamadaani, 2016](#); ([Abed Alrikaby et al., 2020](#), [Alrikaby et al., 2020](#) (in Thailand [Butboonchoo, et al., 2016](#)). in India ([Roy et al., 2008](#)).

Aporina delafondi ([Railliet, 1892](#))

Phylum: Platyhelminthes ([Gegenbaur, 1859](#))
Class: Cestoda ([Rudolphi, 1808](#))
Order: Cyclophyllidea ([Braun, 1900](#))
Family Anoplocephalidae
Intermediate Distribution (In Iraq [Al-Bayati, 2011](#)).

Hymenolepis cantaniana

Phylum: Platyhelminthes ([Gegenbaur, 1859](#))
Class: Cestoda ([Rudolphi, 1808](#))

Order: Hymenolepididae
Cyclophyllidea (Braun, 1900)

Family
Intermediate hosts dung beetle
Distribution: (Nigeria Adang et al., 2008) (Mohammed et al., 2019).
Habitat: small intestine Duodenum

Hymenolepis carioca

Intermediate hosts beetle
Distribution: (Nigeria Adang et al., 2008) (Mohammed et al., 2019).
Habitat: small intestine Duodenum

Amoebotaenia cuneata

Phylum: Platyhelminthes (Gegenbaur, 1859)
Class: Cestoda (Rudolphi, 1808)
Order: Cyclophyllidea (Braun, 1900)
Family: Dilepididae (Railliet & Henry, 1909)
species: *Amoebotaenia cuneata* (von Linstow, 1872)
Intermediate hosts earthworms (e.g. of the genus *Lumbricus*
Distribution: (Nigeria Adang et al., 2008) (Biswal et al., 2014) (Mohammed et al., 2019).
Habitat: small intestine Duodenum

***Davainea proglottina* (Mohammed et al., 2019).**

Phylum: Platyhelminthes (Gegenbaur, 1859)
Class: Cestoda (Rudolphi, 1808)
Order: Cyclophyllidea (Braun, 1900)
Family Davaineidae (Braun, 1900)
Intermediate hosts ants

Conclusion

Based on analysis of literature data ,it was revealed that have 6 geneses of Cestoda in *Columba livia* and they are *Cotugnia*, *Raillietinia Aporina*, *Hymenolepis*, *Amoebotaenia* the genus Genus: *Cotugnia* Diamare, 1893 have 7 species which are *Cotugnia digonopora*, *Cotugnia intermedia*, *Cotugnia polycantha*, *Cotugnia cuneata*, *Cotugnia satpuliensis*, *Cotugnia srivastavai*, *Cotugnia majdoubii* and the genus *Raillietina* Fuhrmann, 1920 have 7 species *Raillietinia canabia*, *Raillietinia zahratis*, *Raillietina tetragona*, *Raillietina cesticillus*, *Raillietina micracantha*, *Raillietina vinagoi* *Raillietina echinobothrida*, *Raillietina magninumida*.

The genus *Aporina* have one species *Aporina delafondi* (Railliet, 1892) and the genus *Hymenolepis* have *Hymenolepis cantaniana* *Hymenolepis carioca* and the genus *Amoebotaenia* have one species which is *Amoebotaenia cuneata* and the genus *Davainea* have one genus *Davainea proglottina* (Mohammed et al., 2019).

Conflict of Interest

The author hereby declares no conflict of interest.

Consent for publication

The author declares that the work has consent for publication.

References

- Abed Alrikaby, N. J., Maktoof, A. A., & Hafedh, A. A. (2020). Bioaccumulation of some heavy elements in different tissue of *Cotugnia polycantha* and two parasites (*Raillietina tetragona* and *Streptopelia senegalensis*) infected with birds. *EurAsian Journal of BioSciences*, 14(1), 395-399.
- Adang, K. L., Oniye, S. L., Ajanusi, O. J., Ezealor, A. U., & Abdu, P. A. (2008). Gastrointestinal helminths of the domestic pigeons (*Columba livia domestica* gmelin, 1789 aves: columbidae) in Zaria, northern Nigeria. *Science World Journal*, 3(1).
- Ahmed, S. E. (2012). Molecular characterization of *Cotugnia polycantha* (Cestoda, Cyclophyllidae, Davaineidae) infecting doves (*Streptopelia senegalensis*) and pigeons (*Columba livia Domestica*) from Egypt. *Life Science Journal*, 1, 9.
- Al-Bayati, N. Y. (2011). A study on pigeons (*Columba livia*) Cestodes infection in Diyala Province. *Diyala Agricultural Sciences Journal*, 3(2), 1-12.
- Al-Hussaini, M. G., & Maktoof, A. A. (2021). Estimation of Lead, Copper and Zinc Elements in Livers and Kidneys of *Columba livia domestica* which infected with *Raillietina* sp. in Al-Nassiriyah City/South of Iraq. *University of Thi-Qar Journal of Science*, 8(1), 67-71.
- Al-Marsomy, W. A., & Al-Hamadaani, H. S. (2016). Association of cestoda *Raillietina echinobothrida* in rock pigeon *Columba livia* from Baghdad city of Iraq. *Baghdad Science Journal*, 13(3), 354-268.
- Alrikaby, N. J., Hafedh, A. A., & Abul-Doanej, H. A. I. (2020). Prevalence of Cestoda in Domestic Pigeons (*Columba Livia*) of Nasiriyah City, Southern of Iraq. *The Journal of Research on the Lepidoptera*, 51(1), 333-338.
- Biswal, D., Nandi, A. P., & Chatterjee, S. (2014). Biochemical and molecular characterization of the Cyclophyllidean cestode, *Cotugnia cuneata* (Meggit, 1924), an endoparasite of domestic pigeons, *Columba livia domestica*. *Journal of parasitic diseases*, 38, 106-110.
- Butboonchoo, P., Wongsawad, C., Rojanapaibul, A., & Chai, J. Y. (2016). Morphology and molecular phylogeny of *Raillietina* spp. (Cestoda: Cyclophyllidae: Davaineidae) from domestic chickens in Thailand. *The Korean Journal of Parasitology*, 54(6), 777.
- Caira, J. N., Jensen, K., Waeschenbach, A., Olson, P. D., & Littlewood, D. T. J. (2014). Orders out of chaos—molecular phylogenetics reveals the complexity of shark and stingray tapeworm relationships. *International Journal for Parasitology*, 44(1), 55-73.
- Diakou, A., Ptochos, S., & Papadopoulos, E. (2013). Cestode fauna of feral pigeons in Thessaloniki; Northern Greece. *Helminthologia*, 50, 39-42.
- Dranzoa, C., Ocaido, M., & Katete, P. (1999). The ecto, gastro-intestinal and haemo-parasites of live pigeons (*Columba livia*) in Kampala, Uganda. *Avian pathology*, 28(2), 119-124.
- Ghobashy, M. A., & Taeleb, A. A. (2015). Molecular characterization of *Raillietina* (R.) spp. Ortlepp, 1938 (Cestoda: Cyclophyllidae: Davaineidae) infecting domestic and wild birds (*Columba livia* and *Columba livia domestica*). *World Journal of Zoology*, 10(2), 136-141.
- Hassan, Z. (2013). The Pathological effect of cestoda *Raillietina cesticillus* in small intestine of local domestic pigeons *Columba livia domesticus* in Basrah Province. *AL-Qadisiyah Journal of Veterinary Medicine Sciences*, 12(1), 21-26.
- Khalil, L. F., Jones, A., & Bray, R. A. (1994). Keys to the cestode parasites of vertebrates. *CABI, Wallingford*.
- Magzoub, M., Kasim, A. A., & Shawa, Y. (1980). Three new species (Cestoda: Davaineidae) from the rock pigeon *Columba livia domestica* with comments on the infection. *Journal of the College of Science, University of Riyadh*, 11, 119-127.
- Malhotra, S. K., & Kapoor, V. N. (1983). A new cestode *Cotugnia satpuliensis* sp. n. from *Columba livia domestica* and *Columba livia intermedia* from India. *Acta Parasitologica Polonica*, 28(38-52).
- Malviya, H. C., & Dutt, S. C. (1970). Morphology and life history of *Cotugnia srivastavai* n. sp. (Cestoda: Davaineidae) from the domestic pigeon. *Morphology and life history of Cotugnia srivastavai n. sp. (Cestoda: Davaineidae) from the domestic pigeon*.

- Marques, S. M., Quadros, R. M., Da-Silva, C. J. and Baldo, M. (2007). Parasites of pigeons (*Columba livia*) in urban areas of lages, Southern Brazil. *Parasitología latinoamericana*, 62(3-4), 183-187.
- Mohammed, B. R., Simon, M. K., Agbede, R. I. S., & Arzai, A. H. (2019). Prevalence of intestinal helminth parasites of pigeons (*Columba livia domestica* Gmelin 1789) in Kano State, North-Western Nigeria. *Veterinary Parasitology: Regional Studies and Reports*, 16, 100289.
- Musa, S., Afroz, S. D., & Khanum, H. (2011). Occurrence of ecto-and endo parasites in pigeon (*Columba livia* Linn.). *University Journal of Zoology, Rajshahi University*, 30, 73-75.
- Ptochos, S., Diakou, A. & Papadopoulos, E. (2013). Cestode fauna of feral pigeons in Thessaloniki; Northern Greece. *Helminthologia*, 50, 39-42.
- Radfar, M. H., Asl, E. N., Seghinsara, H. R., Dehaghi, M. M., & Fathi, S. (2012). Biodiversity and prevalence of parasites of domestic pigeons (*Columba livia domestica*) in a selected semiarid zone of South Khorasan, Iran. *Tropical animal health and production*, 44, 225-229.
- Roy, B., Lalchhandama, K., & Dutta, B. (2008). Scanning electron microscopic observations on the in vitro anthelmintic effects of *Millettia pachycarpa* on *Raillietina echinobothrida*. *Pharmacognosy Magazine*, 4(13), 20.
- Smyth, J. D., & McManus, D. P. (1989). *The physiology and biochemistry of cestodes*. Cambridge university press.
- Torres, J., Foronda, P., Eira, C., Miquel, J., & Feliu, C. (2010). Trace element concentrations in *Raillietina micracantha* in comparison to its definitive host, the feral pigeon *Columba livia* in Santa Cruz de Tenerife (Canary Archipelago, Spain). *Archives of environmental contamination and toxicology*, 58, 176-182.
- Waghmare, S. B., Padwal, N. D., & Jadhav, B. V. (2009). Effect of deproteinizing agents on biochemical variability of total protein in *Cotugnia digonopora*. *International Journal of Pharmaceuticals Analysis*, 1(1), 1.

How to cite this article

Alrikaby, N. J. and Abul-Doanej, H. A. I. (2023). A systematic review of tapeworms (Plathelminthes, Cestoda) of *Columba livia*. *Science Archives*, Vol. 4(2), 102-105. <https://doi.org/10.47587/SA.2023.4206>

This work is licensed under a [Creative Commons Attribution 4.0 International License](#)



Publisher's Note: MD International Publishing stays neutral about jurisdictional claims in published maps and institutional affiliations.