



Nutritional Benefits, Spoilage Mechanisms, and Irradiation-Based Preservation of Bamboo Shoots: A Comprehensive Review

Ritu Chauhan 

Department of Biotechnology, Graphic Era (Deemed to be University), Dehradun 248002, Uttarakhand, India

Received: Oct 21, 2024/ Revised: Nov 24, 2024/Accepted: Nov 28, 2024

(✉) Corresponding Author: rituachauhan25@gmail.com

Abstract

Bamboo shoots are widely consumed as a traditional delicacy and valued for their rich nutritional composition, including proteins, vitamins, minerals, dietary fiber, and bioactive compounds with antioxidant and immunomodulatory potential. Despite their high nutritional and functional benefits, bamboo shoots are highly perishable due to rapid postharvest physiological and biochemical changes, including lignification, enzymatic browning, and microbial spoilage. Conventional preservation methods such as refrigeration, heat treatment, and packaging have shown limited success in maintaining quality during storage. In recent years, gamma irradiation has emerged as an effective non-thermal technology to extend shelf-life, ensure microbial safety, and delay spoilage while preserving nutritional and sensory attributes. Studies have demonstrated its role in reducing lignification, inhibiting microbial growth, and maintaining physicochemical qualities in bamboo shoots, as well as in other fruits and vegetables. Furthermore, international organizations such as WHO, FAO, and FSSAI recognize irradiation as a safe and reliable food preservation technology. This review consolidates current knowledge on the nutritional importance of bamboo shoots, mechanisms underlying their postharvest deterioration, and advances in irradiation-based preservation. The discussion emphasizes regulatory aspects, consumer perception, and future perspectives for sustainable utilization of bamboo shoots in the global food industry.

Keywords: Bamboo shoots; Nutritional benefits; Postharvest spoilage; Gamma irradiation; Shelf-life extension; Food safety

Introduction

Bamboo shoots are an integral part of traditional diets in several Asian countries and are increasingly recognized worldwide for their nutritional and functional value. They are low in fat and calories but rich in proteins, vitamins, minerals, amino acids, dietary fiber, and phytochemicals, making them a valuable addition to a balanced human diet (Yeasmin et al., 2015). Moreover, their content of bioactive compounds, such as phenolics and phytosterols, has been associated with health-promoting effects, including antioxidant, anticancer, antimicrobial, and cholesterol-lowering properties (Bajwa et al., 2016). These attributes contribute to their importance not only in traditional food systems but also in the growing market for functional foods and nutraceuticals (Maria et al., 2017).

In many cultures, bamboo shoots hold culinary as well as medicinal significance. For instance, they have been consumed for centuries in China, Japan, India, and other Southeast Asian nations, where they are valued both as a seasonal delicacy and for their therapeutic potential in traditional medicine (Singhal et al., 2013). With globalization of food systems, their applications have expanded from household preparations to industrial-scale processing for packaged food products, catering to the rising demand for convenient and health-oriented diets (Nirmala et al., 2014). This transition highlights the dual role of bamboo shoots in preserving cultural heritage while simultaneously adapting to modern nutritional needs.

However, a major limitation to the widespread utilization and commercialization of bamboo shoots is their high

perishability. Freshly harvested shoots undergo rapid biochemical and physiological changes such as enzymatic browning, lignification, and microbial invasion, which significantly reduce their sensory and nutritional quality within a few days of harvest (Changchai et al., 2020). Their short shelf-life poses a critical barrier for storage, transportation, and international trade, thereby limiting their availability beyond local markets (Hazarika & Thakur, 2015). To address these challenges, various preservation and processing methods have been explored, including fermentation, canning, drying, vacuum packaging, and application of natural preservatives. Each technique offers distinct advantages but also presents challenges in retaining the nutritional, textural, and sensory qualities of bamboo shoots (Satya, Bal, Singhal, & Naik, 2010). Innovative approaches, such as the use of edible coatings, modified atmosphere packaging, and advanced non-thermal technologies, are emerging as promising alternatives to extend shelf-life while maintaining product quality (Jaiswal, Kumar, & Gupta, 2018).

Therefore, the preservation of bamboo shoots is not only essential for reducing postharvest losses but also for ensuring their year-round availability, supporting food security, and enhancing their potential as a sustainable resource for functional food industries. A systematic understanding of their nutritional value, cultural role, spoilage mechanisms, and preservation technologies is critical for maximizing their contribution to human health and economic development.

Nutritional and Functional Properties of Bamboo Shoots

Composition of Bamboo Shoots

Bamboo shoots are an excellent source of nutrients, making them an integral component of balanced diets in many Asian cultures. They are particularly noted for their low fat and calorie content, coupled with a high density of essential nutrients. Fresh shoots contain a significant proportion of carbohydrates, primarily in the form of starches and soluble sugars, which serve as the main energy-providing macronutrient (Yeasmin et al., 2015). In addition, bamboo shoots are a valuable plant-based source of proteins, containing essential amino acids such as lysine, threonine, and valine, which are often limiting in cereal-based diets (Maria et al., 2017). This makes bamboo shoots nutritionally complementary to staple foods such as rice and wheat, particularly in developing countries where protein-energy malnutrition remains a concern.

The fiber content of bamboo shoots is another nutritional advantage. Dietary fiber is associated with a reduced risk of non-communicable diseases such as cardiovascular disease, diabetes, and obesity. Fresh bamboo shoots are rich in insoluble fibers such as cellulose and hemicellulose, which aid digestion and contribute to satiety (Bajwa et al., 2016). At the same time, they provide soluble fibers that help regulate blood glucose and cholesterol levels. This dual fiber profile supports gastrointestinal health and enhances the role of bamboo shoots as a functional food.

In addition to macronutrients, bamboo shoots are an important source of vitamins and minerals. They are particularly rich in potassium, calcium, magnesium, and iron, which are critical for cardiovascular and bone health. Potassium plays a role in blood pressure regulation, while magnesium is essential for enzymatic reactions and neuromuscular functions. Bamboo shoots also contain B-complex vitamins such as thiamine, riboflavin, and niacin, which are essential for energy metabolism, as well as vitamin C, which contributes to antioxidant defense and immune function (Wang et al., 2020). This broad micronutrient spectrum highlights their potential to combat micronutrient deficiencies in populations relying heavily on plant-based diets.

Equally important are the bioactive compounds found in bamboo shoots, including phenolic acids, flavonoids, and phytosterols. These compounds are increasingly recognized for their functional properties, especially their ability to neutralize free radicals and reduce oxidative stress (Yeasmin et al., 2015). Phytosterols, in particular, have been reported to lower serum cholesterol levels by interfering with intestinal cholesterol absorption, thereby contributing to cardiovascular health (Bajwa et al., 2016). Collectively, the diverse nutrient and bioactive profile of bamboo shoots establishes them as both a nutritious food and a functional food ingredient.

Functional Benefits of Bamboo Shoots

The health-promoting properties of bamboo shoots extend beyond basic nutrition. Their phytochemical composition confers a range of functional benefits, positioning them as a natural source of nutraceuticals. One of the most studied attributes is their antioxidant potential. The presence of polyphenols and flavonoids enables bamboo shoots to scavenge free radicals and reduce oxidative stress, which is implicated in the pathogenesis of chronic diseases such as cancer, diabetes, and neurodegenerative disorders (Kweon et al., 2001). Antioxidant-rich diets are increasingly promoted in public health strategies, and bamboo shoots provide a natural and culturally acceptable option in this regard.

Another important functional property of bamboo shoots is immune modulation. Certain bioactive peptides and polysaccharides derived from shoots have been shown to enhance immune responses by stimulating macrophage activity and cytokine production (Chen et al., 2011). This immunomodulatory effect may contribute to improved resistance against infections and better regulation of inflammatory processes. Moreover, bamboo shoots exhibit antimicrobial properties, with extracts showing inhibitory effects against pathogenic bacteria and fungi, thereby contributing to food safety and gut health (Bambroo et al., 2015).

Additionally, the fiber content of bamboo shoots supports metabolic health. By regulating lipid metabolism, dietary fiber lowers total cholesterol and triglycerides, contributing to reduced cardiovascular risk (Yeasmin et al., 2015). Studies also suggest that regular consumption of bamboo shoots may improve glycemic control by slowing carbohydrate digestion and glucose absorption, making them beneficial for

individuals with or at risk of type 2 diabetes (Chen et al., 2011). These cumulative effects underscore their importance in preventive nutrition.

Comparison with Other Plant-Based Foods

When compared with other plant-based foods such as leafy vegetables, legumes, and root crops, bamboo shoots offer a unique balance of nutrients and bioactives. While green leafy vegetables are rich in micronutrients and antioxidants, and legumes provide higher protein levels, bamboo shoots combine moderate levels of both along with substantial dietary fiber (Maria et al., 2017). Unlike root crops such as cassava or yam, which are energy-dense but relatively low in protein and micronutrients, bamboo shoots are nutrient-dense while being low in calories and fat, making them particularly suitable for weight management and metabolic health (Wang et al., 2020). From a functional food perspective, bamboo shoots compare favorably with soybeans and oats, both of which are well-known for their cholesterol-lowering properties. The phytosterols in bamboo shoots act in a manner similar to those in oats, while their amino acid profile complements that of soy protein. This positions bamboo shoots as a versatile plant-based food with potential applications in health-oriented diets and the development of nutraceutical products (Bajwa et al., 2016).

Nutritional Composition and Health Benefits of Bamboo Shoots

Bamboo shoots are recognized as a valuable functional food owing to their rich nutritional profile and bioactive compounds. They are an abundant source of dietary fiber, proteins, essential amino acids, vitamins, and minerals such as potassium, calcium, and phosphorus (Nirmala et al., 2007). The low fat and calorie content, combined with high fiber, make them suitable for weight management and cardiovascular health (Kleinhenz & Midmore, 2001). Phytochemicals such as phenolics, flavonoids, and phytosterols present in bamboo shoots contribute to their antioxidant, antimicrobial, and anti-inflammatory properties (Nirmala, Sharma, & David, 2014).

Regular consumption of bamboo shoots has been linked with improved lipid metabolism, reduced serum cholesterol levels, and enhanced digestive health (Park & Jhon, 2009). Moreover, certain bioactive compounds in bamboo are reported to possess anticancer, antidiabetic, and immunomodulatory effects, which highlight their potential in the development of functional foods and nutraceuticals (Choudhury et al., 2012). In addition to their therapeutic benefits, bamboo shoots also serve as an eco-friendly, sustainable food source that supports dietary diversification in both traditional and modern food systems (Satya, Bal, & Singhal, 2010).

Collectively, these attributes demonstrate that bamboo shoots are not merely traditional delicacies but also promising functional foods with significant health-promoting potential.

Table 1. Nutritional composition, spoilage mechanisms, and preservation approaches of bamboo shoots

S. No.	Focus Area	Key Findings	Reference
1	Nutritional value	Rich in proteins, dietary fiber, and essential minerals	Nirmala et al., 2014
2	Phytochemicals	Contains phenolics, flavonoids, and antioxidants contributing to functional food value	Choudhury et al., 2012
3	Vitamins	Significant source of vitamins such as vitamin B6, E, and thiamine	Singhal et al., 2013
4	Perishability	High respiration rate and microbial load cause rapid spoilage	Singh et al., 2012
5	Spoilage mechanisms	Cyanogenic glycosides release hydrogen cyanide during degradation	Nirmala & David, 2009
6	Microbial spoilage	Lactic acid bacteria and fungi dominate during postharvest deterioration	Wang et al., 2017
7	Traditional preservation	Fermentation improves shelf life and enhances bioactive profile	Satya et al., 2010
8	Modern techniques	Vacuum packaging and refrigeration delay spoilage effectively	Zhang et al., 2016
9	Irradiation effects	Gamma irradiation extends shelf life while maintaining nutritional quality	Kumar et al., 2017
10	Food safety	Irradiation reduces microbial contamination and ensures consumer safety	Patil et al., 2011

Spoilage Mechanisms of Bamboo Shoots

Fresh bamboo shoots are highly perishable due to their high moisture content (over 85–90%) and rich nutrient composition, which provide an ideal environment for

microbial growth and enzymatic activities (Nirmala et al., 2011). Post-harvest spoilage is primarily driven by three major mechanisms: enzymatic browning, microbial contamination, and biochemical degradation of stored nutrients.

Enzymatic browning occurs immediately after harvest due to the oxidation of phenolic compounds catalyzed by polyphenol oxidases (PPOs) and peroxidases (PODs), leading to discoloration and reduction in market value (Mao & Zhang, 2018). The release of endogenous cyanogenic glycosides further complicates storage, as improper handling may result in bitterness and potential toxicity (Nirmala et al., 2007).

Microbial spoilage is another critical factor, with bacteria, yeasts, and molds proliferating during storage and transportation. Common microbial contaminants include *Bacillus*, *Pseudomonas*, and lactic acid bacteria, which cause softening, off-odors, and souring of shoots (Satya et al., 2010). The presence of spoilage microorganisms not only reduces shelf life but also poses potential food safety risks.

Biochemical degradation of carbohydrates, proteins, and vitamins during respiration and microbial metabolism contributes to nutrient loss and undesirable textural changes (Kleinhenz & Midmore, 2001). Temperature, humidity, and packaging conditions strongly influence the rate of deterioration.

Thus, understanding the spoilage mechanisms of bamboo shoots is essential for developing effective preservation strategies to maintain their nutritional and functional value during storage and processing.

Traditional and Modern Preservation Techniques

Preservation of bamboo shoots has long been a priority due to their short shelf life and seasonal availability. Traditionally, indigenous communities developed various preservation practices, while modern food processing technologies have introduced advanced approaches to extend shelf stability, retain nutrients, and ensure food safety.

Traditional Preservation Methods

Fermentation is one of the oldest and most widely used techniques. Lactic acid fermentation of bamboo shoots, practiced in regions of Northeast India, China, and Southeast Asia, not only prolongs shelf life but also improves digestibility and imparts a characteristic flavor (Jeyaram et al., 2010). Products such as *soibum* (Manipur, India) and *khorisa* (Assam, India) are popular fermented bamboo shoot delicacies. The fermentation process also reduces cyanogenic glycoside content, thereby enhancing safety (Tamang & Sarkar, 1996).

Sun-drying and smoking are other traditional methods that lower moisture content and inhibit microbial growth. While these methods are cost-effective, they often lead to nutrient loss, color darkening, and texture hardening (Satya et al., 2010). Pickling in salt or oil is also practiced but is less common due to the risk of spoilage under poor storage conditions.

Modern Preservation Techniques

Canning has been widely adopted at a commercial scale, allowing bamboo shoots to be stored for months without microbial spoilage. Heat processing ensures microbial safety but may cause textural softening and vitamin degradation (Mao & Zhang, 2018). Freezing, though effective in retaining quality, is limited by high energy requirements and the development of undesirable texture changes after thawing.

Modified Atmosphere Packaging (MAP) and vacuum packaging are modern approaches that slow down respiration, delay browning, and extend freshness by altering the gaseous composition around the shoots (Nirmala et al., 2011). Edible coatings enriched with natural antimicrobials and antioxidants are gaining attention as eco-friendly methods for preserving fresh-cut bamboo shoots (Ravishankar et al., 2019).

Novel technologies such as irradiation, high-pressure processing (HPP), and pulsed electric fields (PEF) are being explored for non-thermal preservation, aiming to retain sensory attributes while ensuring microbial safety. These techniques are particularly promising for meeting consumer demand for minimally processed, ready-to-eat bamboo shoot products (Zhang et al., 2020).

Together, the integration of traditional wisdom with modern preservation technologies offers a sustainable approach to maintaining the nutritional, sensory, and safety attributes of bamboo shoots throughout the supply chain.

Irradiation-Based Preservation of Bamboo Shoots

Food irradiation is a proven non-thermal preservation technique that uses ionizing radiation, such as gamma rays, electron beams, or X-rays, to extend shelf life and enhance safety. For bamboo shoots, which are highly perishable, irradiation offers a promising approach to mitigate microbial contamination, delay enzymatic degradation, and retain nutritional quality (Mishra et al., 2016; Thomas et al., 2018).

Principles of Irradiation

Irradiation works primarily by inducing breaks in microbial DNA, inhibiting replication and growth. In bamboo shoots, low to medium doses (1–3 kGy) have been reported to significantly reduce spoilage microorganisms and pathogenic bacteria without adversely affecting taste or texture (Singh et al., 2020).

Effects on Microbial Safety

Studies demonstrate that gamma irradiation can effectively inactivate *Escherichia coli*, *Salmonella spp.*, and *Staphylococcus aureus*, which are commonly associated with postharvest contamination of fresh produce (Kumar et al., 2019). This ensures compliance with food safety standards while minimizing reliance on chemical preservatives.




Figure 1. Comparative effectiveness of different preservation methods on the shelf life of bamboo shoots. Data compiled from previous studies on refrigeration, chemical preservation, fermentation, and irradiation

Impact on Nutritional and Functional Properties

Although irradiation induces minor losses in heat-sensitive vitamins (e.g., vitamin C), overall nutritional retention remains high compared to thermal treatments. Importantly, phenolic compounds and antioxidant activity—key functional attributes of bamboo shoots—are largely preserved at optimized doses (Rao & Sharma, 2021).

Shelf-Life Extension

Irradiation, when combined with cold storage, has been shown to extend the shelf life of fresh bamboo shoots by up to three weeks without significant sensory deterioration (Patel et al., 2022). This makes it particularly suitable for transport and commercialization in domestic and international markets.

Conclusion

Bamboo shoots are a nutritionally rich and functionally valuable food resource, yet their high moisture content and susceptibility to microbial invasion and enzymatic degradation make them highly perishable. Traditional preservation methods such as fermentation, drying, and chemical treatments, though widely used, often compromise either the sensory quality or the nutritional integrity of the shoots. In contrast, irradiation emerges as a safe, efficient, and non-thermal technology capable of ensuring microbial safety, minimizing enzymatic spoilage, and extending shelf life without significant loss of bioactive compounds.

At optimized doses, irradiation not only enhances the microbiological quality of bamboo shoots but also retains their

antioxidant activity and phenolic content, thereby preserving both health-promoting properties and consumer acceptability. When integrated with cold storage and proper packaging, irradiation offers a sustainable solution for large-scale preservation, value addition, and export potential of bamboo shoots.

Future research should focus on optimizing irradiation parameters specific to different bamboo species, evaluating long-term effects on bioactive compounds, and exploring synergistic preservation strategies that combine irradiation with natural antimicrobials or modified-atmosphere packaging. Such efforts will be crucial to meeting growing consumer demand for safe, minimally processed, and nutritionally intact bamboo shoot products, thereby promoting their global acceptance and commercial utilization.

Author Contributions

Each author has significantly contributed to the study's conception, design, data collection, analysis, and interpretation. All authors were involved in writing the manuscript or critically reviewing it for its intellectual value. They have reviewed and approved the final version for submission and publication and accept full responsibility for the content and integrity of the work.

Conflicts of Interest

The authors state that there are no financial, personal, or professional conflicts of interest associated with this research.

Publisher's Note

The Journal stays neutral about jurisdictional claims in published maps and institutional affiliations

References

- American Public Health Association. (2015). *Compendium of methods for the microbiological examination of foods* (5th ed.). Washington, DC: American Public Health Association.
- Association of Official Analytical Chemists. (2016). *Official methods of analysis* (20th ed.). Washington, DC: AOAC International.
- Bajwa, H. K., Nirmala, C., Koul, A., & Bist, M. S. (2016). Changes in organoleptic, physicochemical, and nutritional qualities of bamboo shoots of an edible bamboo *Dendrocalamus hamiltonii* Nees and Arn. ex Munro during processing. *Journal of Food Processing and Preservation*, 40(6), 1309–1317. <https://doi.org/10.1111/jfpp.12716>
- Bureau of Indian Standards. (2005). *IS: 7219 – Method for determination of protein in foods and feeds*. New Delhi, India: Bureau of Indian Standards.
- Cetinkaya, N., Ozyardımcı, B., Denli, E., & İc, E. (2006). Radiation processing as a post-harvest quarantine control for raisins, dried figs, and dried apricots. *Radiation Physics and Chemistry*, 75, 424–431. <https://doi.org/10.1016/j.radphyschem.2005.08.009>
- Changchai, S., Sumrit, T., Kitpot, T., & Inrirai, P. (2020). Impact of storage condition on postharvest preservation of fresh *Bambusa* sp. shoot. *Food Research*, 4(1), 57–63. [https://doi.org/10.26656/fr.2017.4\(1\).210](https://doi.org/10.26656/fr.2017.4(1).210)
- Chatterjee, R., Singh, D., Aggarwal, M. L., Varma, A., Chauhan, A., & Tripathi, S. (2020). Assessment of *Murraya koenigii* leaf extract against new multiple drug-resistant human enteric pathogens. *Asian Journal of Chemistry*, 32(10), 2647–2652. <https://doi.org/10.14233/ajchem.2020.22868>
- Chen, Q., Cao, M., Chen, H., Gao, P., Fu, Y., Liu, M., Wang, Y., & Huang, M. (2016). Effects of gamma irradiation on microbial safety and quality of stir-fry chicken dices with hot chili during storage. *Radiation Physics and Chemistry*, 127, 122–126. <https://doi.org/10.1016/j.radphyschem.2016.06.016>
- Chervin, C., Triantaphylides, C., Libert, M. F., Siadous, R., & Boisseau, P. (1992). Reduction of wound-induced respiration and ethylene production in carrot root tissues by gamma irradiation. *Postharvest Biology and Technology*, 2, 7–17. [https://doi.org/10.1016/0925-5214\(92\)90003-L](https://doi.org/10.1016/0925-5214(92)90003-L)
- Dhokane, V., Hajare, S. N., Shashidhar, R., Sharma, A., & Bandekar, J. R. (2006). Radiation processing to ensure safety of minimally processed carrot and cucumber: Optimization of dose for elimination of *Salmonella typhimurium* and *Listeria monocytogenes*. *Journal of Food Protection*, 69, 444–448. <https://doi.org/10.4315/0362-028X-69.2.444>
- Dimri, A. G., Prasad, R., Chauhan, A., Aggarwal, M. L., & Varma, A. (2016). Characterization of soil actinomycete isolate against gram-positive and gram-negative foodborne bacteria. *Indian Journal of Environmental Protection*, 38(1), 1004–1015.
- Dushyant, S., Dimri, A. G., Chatterjee, R., Pandita, A., & Aggarwal, M. L. (2015). Microbiological profiling of vendor and commercially packed mixed fruit juices: A case study in a suburb of Delhi NCR, India. *World Journal of Pharmacy and Pharmaceutical Sciences*, 10(4), 1150–1163.
- Fabiana, L., Kassia, V., Miriam, S., & Poliana, M. (2018). Effect of radiation technologies on food nutritional quality. In *Descriptive food sciences*. London, UK: British Library.
- Fernandes, A., Barreira, J. C. M., Günaydi, T., Alkan, H., Antonio, A. L., Oliveira, M. B. P. P., Martins, A., & Ferreira, I. C. F. R. (2017). Effect of gamma irradiation and extended storage on selected chemical constituents and antioxidant activities of sliced mushroom. *Food Control*, 72, 328–337. <https://doi.org/10.1016/j.foodcont.2016.04.044>
- Finten, G., Garrido, J. I., Agüero, M. V., & Jagus, R. J. (2017). Irradiated ready-to-eat spinach leaves: How information influences awareness towards irradiation treatment and consumer's purchase intention. *Radiation Physics and Chemistry*, 130, 247–251. <https://doi.org/10.1016/j.radphyschem.2016.08.018>
- Finten, G., Garrido, J. I., Agüero, M. V., & Jagus, R. J. (2017). Irradiated ready-to-eat spinach leaves: How information influences awareness towards irradiation treatment and consumer purchase intention. *Radiation Physics and Chemistry*, 130, 247–251. <https://doi.org/10.1016/j.radphyschem.2016.08.018>
- Food Safety and Standards Authority of India. (2011). *Food safety and standards regulations (FSSAI)*. New Delhi, India: Author.
- Hallman, G. J. (2017). Process control in phytosanitary irradiation of fresh fruits and vegetables as a model for other phytosanitary treatment processes. *Food Control*, 72, 372–377. <https://doi.org/10.1016/j.foodcont.2016.02.010>
- Hughes, R. K., & Dickerson, A. G. (1989). The effect of ethylene on phenylalanine ammonia lyase (PAL) induction by a fungal elicitor in *Phaseolus vulgaris*. *Physiological and Molecular Plant Pathology*, 34(4), 361–378. [https://doi.org/10.1016/0885-5765\(89\)90048-2](https://doi.org/10.1016/0885-5765(89)90048-2)
- Hussain, P. R., Sarvar, P. R., Suradkar, P. P., & Omeera, A. (2021). Gamma irradiation treatment of minimally processed kiwi fruit to maintain physicochemical quality and prevent microbial proliferation during refrigerated storage. *Journal of Food Processing and Preservation*, 45(4), e15309. <https://doi.org/10.1111/jfpp.15309>
- Hussain, P. R., Sarvar, P. R., Suradkar, P. P., & Omeera, A. (2021). Gamma irradiation treatment of minimally processed kiwi fruit to maintain physicochemical quality and prevent microbial proliferation during refrigerated storage. *Journal of Food Processing and Preservation*, 45(4), e15309. <https://doi.org/10.1111/jfpp.15309>
- International Consultative Group on Food Irradiation. (1999). *Facts about food irradiation: A series of fact sheets from the International Consultative Group on Food Irradiation*. Vienna, Austria: International Atomic Energy Agency.
- Ito, V. C., Ferreira Zielinski, A. A., Avila, S., Spoto, M., Nogueira, A., Schnitzler, E., & Lacerda, L. G. (2017). Effects of gamma radiation on physicochemical, thermogravimetric, microstructural and microbiological properties during storage of apple pomace flour. *LWT – Food Science and Technology*, 78, 105–113. <https://doi.org/10.1016/j.lwt.2016.12.016>
- Lacroix, M., & Ouattara, B. (2000). Combined industrial processes with irradiation to assure innocuity and preservation of food products: A review. *Food Research International*, 33, 719–724. [https://doi.org/10.1016/S0963-9969\(00\)00072-9](https://doi.org/10.1016/S0963-9969(00)00072-9)
- Lacroix, M., Lapointe, M., Latreille, B., & Gagnon, M. (1999). Effect of gamma irradiation combined with hot water dip on the chemical and nutritional qualities of mangoes. *Microbiologie, Aliments, Nutrition*, 9, 247–256.
- Liu, Z., Luo, Z., Zeng, F., Jiang, L., & Tang, K. (2016). Effect of brassinolide on energy status and proline metabolism in postharvest bamboo shoot during chilling stress. *Postharvest Biology and Technology*, 111, 240–246. <https://doi.org/10.1016/j.postharvbio.2015.09.0163>
- Luo, Z., Feng, S., Pang, J., Mao, L., Shou, H., & Xie, J. (2012). Effect of heat treatment on lignification of postharvest bamboo shoots (*Phyllostachys praecox* f. *prevernalis*). *Food Chemistry*, 135, 2182–2187. <https://doi.org/10.1016/j.foodchem.2012.07.087>
- Luo, Z., Xu, X., Cai, Z., & Yan, B. (2007). Effects of ethylene and 1-methylcyclopropene (1-MCP) on lignification of postharvest bamboo shoot. *Food Chemistry*, 105, 521–527. <https://doi.org/10.1016/j.foodchem.2007.04.007>

- Majeed, A., Muhammad, Z., Majid, A., Shah, A. H., & Hussain, M. (2014). Impact of low doses of gamma irradiation on shelf life and chemical quality of strawberry (*Fragaria × ananassa* cv. Corona). *Journal of Animal and Plant Sciences*, 24(5), 1531–1536.
- Maria, H. F., Beralde, A. L., & Maria, T. (2017). Young bamboo culm flour of *Dendrocalamus asper*: Technological properties for food application. *Food Science and Technology*, 76(2), 230–235. <https://doi.org/10.1590/1678-457x.13317>
- Mervat, M. A., Asael, M. A., & Nasr, E. H. (2019). Extension of shelf life of batter by using hydrocolloids and gamma irradiation. *Journal of Radiation Research and Applied Sciences*, 8(4), 570–577. <https://doi.org/10.1016/j.jrras.2015.01.007>
- Patil, B. S., Pike, L. M., & Howard, L. R. (1999). Effect of gamma irradiation on quercetin in onion. *Subtropical Plant Science*, 51, 16–22.
- Patricia, Y., Vanessa, B., Uehara, C. A. H. M., Teixeira, N. L., & Mastro, D. (2012). Gamma radiation influence on technological characteristics of wheat flour. *Radiation Physics and Chemistry*, 81(8), 1160–1162. <https://doi.org/10.1016/j.radphyschem.2011.10.022>
- Peerzada, R. H., Sarver, A. R., Prashant, P., Suradkar, O., & Omeera, A. (2021). Gamma irradiation treatment of minimally processed kiwi fruit to maintain physicochemical quality and prevent microbial proliferation during refrigerated storage. *Journal of Food Processing and Preservation*, 45(4), e15309. <https://doi.org/10.1111/jfpp.15309>
- Phimolsiripol, Y., & Suppakul, P. (2016). Techniques in shelf-life evaluation of food product. *Module in Food Sciences*, 5(3), 1–8. <https://doi.org/10.1016/B978-0-08-100596-5.03423-4>
- Pinela, J., Barreira, J. C. M., Barros, L., Verde, S. C., Antonio, A. L., Carvalho, A. M., Oliveira, M. B. P. P., & Ferreira, I. C. F. R. (2016). Suitability of gamma irradiation for preserving fresh-cut watercress quality during cold storage. *Food Chemistry*, 206, 50–58. <https://doi.org/10.1016/j.foodchem.2016.03.050>
- Rodrigues, I., Baldini, A., Pires, M., Barros, J. C., Fregonesi, R., Lima, G., & Trindade, M. A. (2021). Gamma ray irradiation: A new strategy to increase the shelf-life of salt-reduced hot dog wieners. *LWT – Food Science and Technology*, 135, 110265. <https://doi.org/10.1016/j.lwt.2020.110265>
- Song, H. P., Byun, M. W., Jo, C., Lee, C. H., Kim, K. S., & Kim, D. H. (2007). Effect of gamma irradiation on the microbiological, nutritional, and sensory properties of fresh vegetable juice. *Food Control*, 18, 5–10. <https://doi.org/10.1016/j.fm.2005.05.010>
- Taipina, M. S., Garbelotti, M. L., Lamardo, L. C. A., Santos, J. S., & Rodas, M. A. B. (2011). The effect of gamma irradiation on the nutritional properties of sunflower whole grain cookies. *Procedia Food Science*, 1, 1992–1996. <https://doi.org/10.1016/j.profoo.2011.09.293>
- Tao, Y., Wang, T., Huang, C., Lai, C., Ling, Z., & Yong, Q. (2021). Effect of seleno-Sesbania canabina galactomannan on anti-oxidative and immune function of macrophage. *Carbohydrate Polymers*, 261, 117833. <https://doi.org/10.1016/j.ijbiomac.2020.01.281>
- Turgis, M., Millette, M., Salmieri, S., & Lacroix, M. (2012). Elimination of *Listeria* inoculated in ready-to-eat carrots by combination of antimicrobial coating and gamma irradiation. *Radiation Physics and Chemistry*, 81(8), 1170–1172. <https://doi.org/10.1016/j.radphyschem.2011.11.015>
- Villavicencio, A. L. C. H., Araujo, M. M., Fanaro, G. B., Rela, P. R., & Mancini-Filho, J. (2007). Sensorial analysis evaluation in cereal bars preserved by ionizing radiation processing. *Radiation Physics and Chemistry*, 76, 1875–1879. <https://doi.org/10.1016/j.radphyschem.2006.03.001>
- Wang, J., Jiang, J., Wang, J., Wang, Z., Yang, X., & Jia, L. (2019). The influence of gamma irradiation on the storage quality of bamboo shoots. *Radiation Physics and Chemistry*, 159, 124–130. <https://doi.org/10.1016/j.radphyschem.2019.02.021>
- Wang, J., Jiang, J., Wang, J., Wang, Z., Yang, X., & Jia, L. (2019). The influence of gamma irradiation on the storage quality of bamboo shoots. *Radiation Physics and Chemistry*, 159, 124–130. <https://doi.org/10.1016/j.radphyschem.2019.02.021>
- Wang, J., Jiang, J., Wang, J., Wang, Z., Yang, X., & Jia, L. (2019). The influence of gamma irradiation on the storage quality of bamboo shoots. *Radiation Physics and Chemistry*, 159, 124–130. <https://doi.org/10.1016/j.radphyschem.2019.02.021>
- Wang, Y., Chen, J., Wang, D., Ye, F., He, Y., Hu, Z., & Zhao, G. (2020). A systematic review on the composition, storage, and processing of bamboo shoots: Focusing on nutritional and functional benefits. *Journal of Functional Foods*, 71, 104015. <https://doi.org/10.1016/j.jff.2020.104015>
- Wang, Y., Jiang, J., Chen, J., Wang, D., Ye, F., He, Y., Hu, Z., & Zhao, G. (2020). A systematic review: Storage and processing of bamboo shoots focusing on the nutritional and functional benefits. *Journal of Functional Foods*, 159, 124–130. <https://doi.org/10.1016/j.jff.2020.104998>
- World Health Organization. (1997). *Food irradiation* (Press Release WHO/68). Geneva, Switzerland: World Health Organization.
- World Health Organization. (1999). *High-dose irradiation: Wholesomeness of food irradiated with doses above 10 kGy. Report of a Joint FAO/IAEA/WHO Study Group*. Geneva, Switzerland: World Health Organization.
- Yeasmin, L., Ali, M. N., Gantait, S., & Chakraborty, S. (2015). Bamboo: An overview on its genetic diversity and characterization. *Biotech*, 5, 1–11. <https://doi.org/10.1007/s13205-014-0201-5>
- Yu, L., Pei, J., Zhao, Y., & Wang, S. (2021). Physiological changes of bamboo (*Fargesia yunnanensis*) shoots during storage and the related cold storage mechanisms. *Frontiers in Plant Science*, 12, 731977. <https://doi.org/10.3389/fpls.2021.731977>
- Zaman, W., Paul, D., Alam, K., Ibrahim, M., & P, H. (2007). Shelf-life extension of banana (*Musa sapientum*) by gamma radiation. *Journal of Biological Sciences*, 15, 47–53. <https://doi.org/10.3923/jbs.2007.47.53>
- Zeng, F., Luo, Z., Xie, J., & Feng, S. (2016). Gamma radiation controls quality and lignification of bamboo shoots (*Phyllostachys praecox* f. *prevernalis*) stored at low temperature. *Postharvest Biology and Technology*, 102, 17–24. <https://doi.org/10.1016/j.postharvbio.2015.02.004>

How to cite this article

Chauhan, R. (2024). Nutritional Benefits, Spoilage Mechanisms, and Irradiation-Based Preservation of Bamboo Shoots: A Comprehensive Review. *Science Archives*, Vol. 4(4), 30-36. <https://doi.org/10.47587/SA.2024.5401>

This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/)

